

Heavy-baryon chiral perturbation theory approach to thermal neutron capture on ^3He

Rimantas Lazauskas,^{1,*} Young-Ho Song,^{2,†} and Tae-Sun Park^{3,4,‡}

¹*IPHC, IN2P3-CNRS/Université Louis Pasteur BP 28,*

F-67037 Strasbourg Cedex 2, France

²*Department of Physics, Duke University, Durham, NC 27708, USA*

³*Department of Physics and BAERI,*

Sungkyunkwan University, Suwon 440-746, Korea

⁴*Department of Physics and Astronomy,*

University of South Carolina, Columbia, SC 29208, USA

(Dated: May 19, 2009)

Abstract

For radiative thermal neutron capture on ^3He ($^3\text{He} + n \rightarrow ^4\text{He} + \gamma$, known as the *hen* reaction), the cross section is calculated based on heavy-baryon chiral perturbation theory. The relevant M1 operators are calculated up to next-to-next-to-next-to-leading order (N³LO). The nuclear wave functions are obtained from the rigorous Faddeev-Yakubovski equations for five sets of the realistic nuclear Hamiltonian. Up to N³LO, the M1 operators contain two low-energy constants which appear as the coefficients of the non-derivative two-nucleon contact terms. After determining these two constants using the experimental values of the magnetic moments of the triton and ^3He , we have made a totally parameter-free calculation of the *hen* cross section. The results are in good agreement with the data.

PACS numbers: 25.40.Lw, 23.20.-g, 25.10.+s, 11.10.Hi

Keywords:

*Electronic address: rimantas.lazauskas@ires.in2p3.fr

†Electronic address: yhsong@phy.duke.edu

‡Electronic address: tspark@kias.re.kr

I. INTRODUCTION

The radiative capture of a slow neutron on ^3He ($^3\text{He} + n \rightarrow ^4\text{He} + \gamma$), or the *hen* process, is an example of rare situations where the contributions of the single-nucleon (1B) currents are strongly suppressed due to the pseudo-orthogonality of the wave functions; viz., since the major components of the initial and final state wave functions belong to different representations of symmetry group, they cannot be connected through the r -independent leading 1B operators. In fact, this suppression is so strong that the meson-exchange-current (MEC) “corrections” become comparable to the 1B contributions, and furthermore, since the MEC and 1B terms in this case come with opposite signs, a strong cancellation between them occurs, leading to a further drastic suppression of the cross section. The *hen* process near threshold is governed by the M1 operators, and the MEC contributions to them consist of the well-known one-pion-exchange part and the short-range part. It is to be noted that the latter is not constrained by the symmetries of QCD, and that because of the above-mentioned suppression and the cancellation mechanisms, the short-range contributions are crucial even for a rough estimation of the *hen* cross section. Furthermore, the strong suppression of the 1B matrix elements implies that their values are highly sensitive to the details of the wave functions. Therefore, for a precise estimation of the *hen* cross section, it is imperative to have: (i) a reliable method for deriving the relevant MEC operators with a good control of short-range physics, and (ii) the accurate wave functions for the initial and final wave functions. These requirements make the quantitative estimation of the *hep* cross section highly non-trivial.

The strong enhancement of the relative importance of MEC in the *hen* process makes it a valuable testing ground for our understanding of the roles of MEC in light nuclei. Apart from this point, which is important on its own right, a careful study of the *hen* process is also of great importance in connection with the so-called *hep* processes, $^3\text{He} + p \rightarrow ^4\text{He} + \nu_e + e^+$, because *hep* shares all the aforementioned features of *hen*: the drastic suppression of the 1B contributions, strong cancellation between the 1B and 2B contributions, and the sensitivity of the transition amplitude to the details of the nuclear wave functions. The *hep* is one of the proton-burning reactions that take place in the interior of the Sun and, since *hep* produces the highest-energy solar neutrinos among the *pp*-chain reactions, it has attracted much attention in the study of the solar neutrinos; see [1, 2] for a recent review. In a

recent paper, Park *et al.* [3] developed an effective field theory (EFT) approach, which has come to be known as “more-effective EFT” (MEEFT for short) [4], and, with the use of MEEFT, they calculated the *hep* S -factor with an estimated accuracy of about 15 %. In view of the fact that the previous theoretical estimations of the *hep* S -factor ranged over two orders of magnitude [3], this is a remarkable feat. A direct test of this theoretical result, however, is not possible because of the extreme difficulty of measuring the *hep* cross section. Meanwhile, the threshold *hen* cross section has been measured with a reasonable accuracy: $\sigma_{exp} = (54 \pm 6) \mu b$ [5] and $\sigma_{exp} = (55 \pm 3) \mu b$ [6]. Given the close similarity between *hen* and *hep*, it is expected that comparison between theory and experiment for the *hen* case will offer valuable information on the validity of the theoretical framework employed for the *hep* calculation in [3]. This gives an additional motivation for carrying out a detailed study of *hen*.

Although the theoretical investigation of *hen* has a long history, the *hen* cross section has never been explained in a satisfactory manner. Towner and Khanna [7] and Wervelman *et al.* [6] performed shell-model calculations and obtained $\sigma = (14 \sim 125) \mu b$ and $\sigma = (47 \pm 18) \mu b$, respectively. Much more elaborate calculations have been performed by Carlson *et al.* [8] and by Schiavilla *et al.* [9], arriving at $\sigma = 112 \mu b$ and $\sigma = 86 \mu b$, respectively. These works are based on the conventional approach, the so-called *standard nuclear physics approach* (SNPA for short), which consists in the use of phenomenological transition operators and phenomenological wave functions. SNPA has been enormously successful in correlating and explaining a vast range of electroweak nuclear transitions in light nuclei but, from a formal point of view, it has an insufficiency that it is not equipped with a systematic way of deriving MEC operators. The MEC operators in SNPA are constructed based on the approach of Chemtob and Rho [10]. Although this construction of the MEC operators is guided by chiral symmetry and the Ward identities related to the adopted Hamiltonians, a certain degree of model dependence is unavoidable.

In this paper we report on a parameter-free MEEFT calculation for the *hen* cross section at threshold which is based on essentially the same method as adopted in [3].^{#1} In MEEFT the transition operators are derived in a systematic expansion based on HBChPT, and the

^{#1} There has been an attempt by Y.H. Song et al. [11] to carry out an MEEFT calculation of *hen*, but an approximate treatment of the scattering wave function in [11] limits its reliability.

nuclear matrix elements are obtained by sandwiching these operators between wave functions generated from a high-precision phenomenological nuclear Hamiltonian. Mismatch in the short-range is overcome by the renormalization procedure for the local operators, according to the premise of low-energy EFTs (see below). It thus takes advantage of the systematic nature of EFTs and the availability of state-of-art wave functions. The mentioned “formal” mismatch may be regarded as a weak point, and also the accurate reproduction of the relevant effective-range parameters (ERPs) is not automatically guaranteed in MEEFT; we shall come back to these points later in the text.

In the present work we derive the M1 operators within HBChPT up to $N^3\text{LO}$. These M1 operators turn out to contain two low-energy constants (LECs), denoted by g_{4s} and g_{4v} , which are the coefficients of two-nucleon contact-term operators. These LECs can be fixed by requiring that the experimental values of the magnetic moments of the triton and ^3He be reproduced; this is the same strategy as adopted in Refs. [12, 13], where the M1 properties of the $A=2$ and $A=3$ systems were studied in MEEFT. A remark is in order here on how the short-range contributions are taken into account in MEEFT. The basic premise of EFT is that physics pertaining to ranges shorter than the experimentally relevant scale is to be lodged in local operators. It means that, provided that a proper renormalization procedure is implemented to the coefficients of local operators (*i.e.*, LECs), the renormalization invariance ensures that the net physical amplitudes be independent of the details of short-range physics. We implement the renormalization condition here by adjusting the values of LECs (g_{4s} and g_{4v} for *hen* and $\hat{d}^R$ for *hep*) so as to reproduce a set of known experimental data [$\mu(^3\text{H})$ and $\mu(^3\text{He})$ for *hen* and the tritium-beta-decay rate for *hep*]. This matching procedure should be done for each cutoff value and for each Hamiltonian adopted.^{#2} Differences in short-range contributions calculated for each case shift the values of LECs (which are not physical observables), but the physical amplitudes should remain unaffected if renormalization invariance is to hold. The validity of this scheme can be checked by monitoring the stability of the relevant physical observables with respect to changes in the cutoff parameter Λ . It turns out (see below) that, in the present *hen* case, the inclusion of the local-operator (or contact term) contributions reduces the Λ -dependence by a factor of ~ 5 , demonstrating the validity of the adopted renormalization procedure. The residual Λ -dependence may be

^{#2} The details of the cutoff parameter Λ and nuclear Hamiltonians used will be given later in the text.

ascribed to higher order contributions.

On the other hand, for A -body systems with $A \geq 3$, even a realistic nuclear Hamiltonian often fails to reproduce accurately the ERPs that govern the long-range part of the transition matrix elements. If such a mismatch in the long-range region happens, it cannot be cured by local operators, which means that one in general cannot expect to have a reliable theoretical prediction. As an exception to this general statement, however, we should mention that, if a strong correlation between the ERPs and the transition amplitude under consideration can be established with high precision, this correlation can be used to drastically reduce the model dependence of the calculated transition amplitude [13]. We shall take advantage of this feature later in the text (Subsection III A) to obtain as the best estimate for the threshold *hen* cross section: $\sigma = (49.4 \pm 8.5) \mu b$ (for the AV18+UIX potential), and $(44.4 \pm 6.7) \mu b$ (for the I-N3LO+UIX potential); see eq.(17). Good agreement of these estimates with the experimental value of the *hen* cross section gives sound support to the previous MEEFT calculation for the *hep* S -factor [3].

We wish to emphasize that the present work is the first calculation of *hen* that employs nuclear wave functions which are fully consistent with a given realistic nuclear Hamiltonian; these wave functions are numerically exact solutions to the Faddeev-Yakubovski equations in configuration space for a specified realistic nuclear Hamiltonian. It is to be noted that hitherto even the most advanced realistic calculations [8, 9, 11] contained some artificial adjustments to suppress the ${}^3\text{He}(n, p){}^3\text{H}$ channel in the scattering wave functions.

This paper is organized as follows. In the next section we explain the formalisms used to derive the M1 operators and to obtain the four-body nuclear wave functions. Section 3 gives the numerical results and analysis. In the final section the implication of our work is discussed.

II. FORMALISM

A. Electromagnetic current and capture cross section

We derive here the electromagnetic (EM) currents and M1 from HBChPT up to N³LO in chiral order counting. HBChPT contains nucleons and pions as pertinent degrees of freedom, with all the other massive fields integrated out. In HBChPT, the EM currents and M1

operators are expanded systematically with increasing powers of Q/Λ_χ , where Q stands for a typical momentum scale of the process and/or the pion mass, and $\Lambda_\chi \sim 4\pi f_\pi \sim m \sim 1$ GeV is the chiral scale, $f_\pi \simeq 92.4$ MeV is the pion decay constant, and m is the nucleon mass. We remark that, while the nucleon momentum $\mathbf{p}_i$ is of the order of Q , its energy ($\sim \mathbf{p}_i^2/m$) is of the order of Q^2/m , and consequently the four-momentum of the emitted photon q^μ should also be counted as $\mathcal{O}(Q^2/m)$.

The M1 operators up to N³LO consist of 1B and 2B contributions, while three-body operators in our counting scheme enter only at N⁴LO or higher orders. In the center-of-mass frame, each current can be written in the form of

$$\mathbf{J}_{em} = e^{-i\mathbf{q}\cdot\mathbf{X}}(i\mathbf{q} \times \mathbf{j}_\mu + \mathbf{j}_c), \quad (1)$$

where $\mathbf{q}$ is the momentum carried out by the photon, $\mathbf{X} = \mathbf{r}_i$ for the 1B current pertaining to the i -th nucleon, while $\mathbf{X} = (\mathbf{r}_j + \mathbf{r}_k)/2$ for the 2B currents involving the nucleons j and k . The resulting M1 operator in multipole expansion reads

$$\boldsymbol{\mu}_{1M}(q) = \frac{1}{\mu_N} \sum_X \left[\frac{3j_1(qX)}{2q} \hat{\mathbf{X}} \times \mathbf{j}_c + j_0(qX) \mathbf{j}_\mu - \frac{j_2(qX)}{2} \left(\mathbf{j}_\mu - \hat{\mathbf{X}} \hat{\mathbf{X}} \cdot \mathbf{j}_\mu \right) \right]_M, \quad (2)$$

where the subscript M is the magnetic quantum number, $q \equiv |\mathbf{q}| = 20.578$ MeV and $j_n(x)$ is the spherical Bessel function; the sum runs over all nucleons for 1B current and over all pairs of nucleons for the 2B current.

In terms of $\boldsymbol{\mu}_{1M}(q)$, the *hen* cross section at thermal energy is given by

$$\sigma = \alpha\pi \frac{c}{v_{rel}} \left(\frac{q}{mc^2} \right)^2 \left(\frac{q}{\hbar c} \right) |\mathcal{M}|^2 \quad (3)$$

with

$$\mathcal{M} \equiv \left\langle \Psi_{4\text{He}}^{J=0, M=0} \right| \boldsymbol{\mu}_{11}(q) \left| \Psi_{n^3\text{He}}^{J=1, M=-1} \right\rangle, \quad (4)$$

where α is the fine structure constant, and $v_{rel} = 2200$ m/s is the velocity of a thermal neutron.

The explicit forms of the M1 operators up to N³LO are given in our recent papers[12, 13], so we present here only a brief summary of them. The 1B M1 operators including the

relativistic corrections read

$$\begin{aligned}
\boldsymbol{\mu}_{1B}(q) = & \sum_i \frac{1}{2m_p} \left\{ \hat{j}_0(qr_i) \left[\boldsymbol{\sigma}_i \left(\mu_i - Q_i \frac{\bar{\mathbf{p}}_i^2}{2m^2} \right) - \frac{\mu_i - Q_i}{2m^2} \bar{\mathbf{p}}_i \boldsymbol{\sigma}_i \cdot \bar{\mathbf{p}}_i \right] \right. \\
& + \hat{j}_1(qr_i) \left[Q_i \mathbf{r}_i \times \bar{\mathbf{p}}_i \left(1 - \frac{\bar{\mathbf{p}}_i^2}{2m^2} \right) - \frac{w(2\mu_i - Q_i)}{4m} i \mathbf{r}_i \times (\bar{\mathbf{p}}_i \times \boldsymbol{\sigma}_i) \right] \\
& \left. + \frac{(qr_i)^2}{30} \hat{j}_2(qr_i) \mu_i (3\hat{\mathbf{r}}_i \hat{\mathbf{r}}_i \cdot \boldsymbol{\sigma}_i - \boldsymbol{\sigma}_i) + \dots \right\} \quad (5)
\end{aligned}$$

where $\omega = q$ is the photon energy, $\hat{j}_n(x) \equiv \frac{(2n+1)!!}{x^n} j_n(x)$, Q_i and μ_i are the charge and magnetic moments of the i -th nucleon, respectively, and $\bar{\mathbf{p}}_i$ is the mean momentum operator of the i -th nucleon.

The two-body M1 operators up to N³LO can be divided into the *soft*-one-pion-exchange (1π) term, the vertex corrections to the one-pion exchange ($1\pi C$) term, the two-pion-exchange (2π) term, and the contact-term contribution (CT),

$$\boldsymbol{\mu}_{2B}(q) = \sum_{i < j} [\boldsymbol{\mu}_{ij}^{1\pi} + (\boldsymbol{\mu}_{ij}^{1\pi C} + \boldsymbol{\mu}_{ij}^{2\pi} + \boldsymbol{\mu}_{ij}^{\text{CT}})] = \text{NLO} + \text{N}^3\text{LO}. \quad (6)$$

In this work we neglect the so-called fixed-current contribution, which is proved to be numerically negligible [12]. In performing Fourier transformation of the 2B operators into coordinate space, we introduce a Gaussian regulator with a cutoff Λ .

It is to be noted that the contact-term $\boldsymbol{\mu}_{ij}^{\text{CT}}$ contains two low-energy constants (LECs), g_{4s} and g_{4v} ,

$$\boldsymbol{\mu}_{ij}^{\text{CT}} = \frac{1}{2m_p} [g_{4s}(\boldsymbol{\sigma}_i + \boldsymbol{\sigma}_j) + g_{4v}(\vec{\tau}_i \times \vec{\tau}_j)^z(\boldsymbol{\sigma}_i \times \boldsymbol{\sigma}_j)] \delta_\Lambda^{(3)}(\mathbf{r}_{ij}). \quad (7)$$

Since the values of LECs are not determined by the symmetry arguments, they need to be fixed either by solving QCD at low-energy or by a set of experimental observables that are sensitive to these LECs. Since the former is not feasible at present, we resort to the latter. Specifically, the values of g_{4s} and g_{4v} are fixed so as to reproduce the experimental values of the triton and ³He magnetic dipole moments, for each nuclear interaction model and for each cutoff value.

B. Faddeev-Yakubovskii equations

Four-nucleon wave functions are obtained by solving the Faddeev-Yakubovskii (FY) equations in configuration space [14, 15]. The FY formalism offers a mathematically rigorous

description for both continuum and bound states. In this formalism wave functions are naturally decomposed into so-called FY amplitudes (FYAs). For $A = 4$ systems, there appear two types of FYAs, which we refer to as components K and H . The asymptotes of the components K and H incorporate 3+1 and 2+2 particle channels, respectively (see Fig.1). The FYAs are evaluated in isospin formalism, *i.e.*, protons and neutrons are regarded as




FIG. 1: The FY components, K and H . As $z \rightarrow \infty$, the K components describe 3+1 particle channels, while the H components contain asymptotic states of 2+2 channels, figure [a]. Figure [b] shows the j-j coupling scheme used in expanding K and H into partial wave bases.

the degenerate states with the same mass, which is fixed to $\hbar^2/m = 41.47 \text{ MeV} \cdot \text{fm}^2$. With the three-body force $V_{123} = V_{12}^3 + V_{23}^1 + V_{31}^2$, the FY equations for $K \equiv K_{1,23}^4$ and $H \equiv H_{12}^{34}$ read [16, 17]

$$\begin{aligned} \left(E - H_0 - V_{12} - \sum_{i < j} V_{ij}^C \right) K &= V_{12}(P^+ + P^-) [(1 + Q)K + H] + \frac{1}{2} (V_{23}^1 + V_{31}^2) \Psi, \\ \left(E - H_0 - V_{12} - \sum_{i < j} V_{ij}^C \right) H &= V_{12} \tilde{P} [(1 + Q)K + H], \end{aligned} \quad (8)$$

where V_{ij} and V_{ij}^C are, respectively, the short-ranged part and the Coulomb-dominated long-range part of the interaction between the i -th and j -th nucleons. $P^+ = (P^-)^{-1} \equiv P_{23}P_{12}$, $Q \equiv -P_{34}$ and $\tilde{P} \equiv P_{13}P_{24} = P_{24}P_{13}$, where P_{ij} is the particle permutation operator. In terms of the FYAs, the total wave function of an $A = 4$ system is given by

$$\Psi = [1 + (1 + P^+ + P^-)Q] (1 + P^+ + P^-)K + (1 + P^+ + P^-)(1 + \tilde{P})H. \quad (9)$$

We expand K and H in terms of the tripolar harmonics $Y_i^\alpha(\hat{x}_i, \hat{y}_i, \hat{z}_i)$, which comprise spins and isospins of the nucleons as well as angular momentum variables,

$$\Phi_i(\vec{x}_i, \vec{y}_i, \vec{z}_i) = \sum_{\alpha} \frac{\mathcal{F}_i^\alpha(x_i, y_i, z_i)}{x_i y_i z_i} Y_i^\alpha(\hat{x}_i, \hat{y}_i, \hat{z}_i), \quad (10)$$

where Φ stands for either K or H , and the subscript i denotes the particle-grouping class (among the four nucleons). We note that the total angular momentum and its projection, parity and the third component of the isospin ($\mathcal{T}_z = 0$) are good quantum numbers, and the subscript α denotes collectively other non-fixed 11 quantum numbers. We use the j - j scheme for the coupling of the angular momenta, as illustrated in Fig. 1b. The Jacobi coordinates used in our study are depicted in Fig. 1. This choice of coordinates allows us to separate the center-of-mass motion and guarantees that kinetic energy operator is independent of the angular variables.

By projecting each of the Eqs. (8) on its natural configuration space basis one obtains coupled integro-differential equations for the radial FYAs ($\mathcal{F}_i^\alpha(x_i, y_i, z_i)$). Note that, contrary to the $3N$ problems, the number of radial FYAs is infinite even when the pair interaction is restricted to a finite number of partial waves. This divergence comes from the existence of the additional degree of freedom l_z in the expansion of the K -type components. Therefore an additional truncation is needed in numerical calculation by taking into account only the relevant amplitudes.

C. Boundary conditions

Eqs. (8) needs to be supplemented with appropriate boundary conditions, which can be written in the Dirichlet form. For both bound and scattering states, the FYAs satisfy the regularity conditions:

$$\mathcal{F}_i^\alpha(0, y_i, z_i) = \mathcal{F}_i^\alpha(x_i, 0, z_i) = \mathcal{F}_i^\alpha(x_i, y_i, 0) = 0. \quad (11)$$

For bound state problems, since the wave functions are compact, the regularity conditions can be completed by forcing the amplitudes $\mathcal{F}_i^\alpha$ to vanish at the borders of the hypercube, $[0, X_{\max}] \times [0, Y_{\max}] \times [0, Z_{\max}]$,

$$\mathcal{F}_i^\alpha(X_{\max}, y_i, z_i) = \mathcal{F}_i^\alpha(x_i, Y_{\max}, z_i) = \mathcal{F}_i^\alpha(x_i, y_i, Z_{\max}) = 0. \quad (12)$$

The hypercube is chosen large enough to accommodate the wave functions.

On the other hand, a scattering state near threshold contains two coupled channels, n - ^3He and p - ^3H , both of which are of type K . For these amplitudes, we impose following matching

condition at $z_i = Z_{\max}$,

$$\begin{aligned} \mathcal{K}_i^\alpha(x_i, y_i, Z_{\max}) &= \frac{1}{\sqrt{4}} \sum_{j'_z l'_z T_3^z} \{f_i^{\alpha a}(\mathbf{x}_i, \mathbf{y}_i)\}_{J_3 \equiv \frac{1}{2}, T_3 T_3^z} \otimes \{Y_{l'_z}(\hat{z}_i) \otimes s_i\}_{j'_z} \rangle_{JM} \\ &\times \left(\frac{i}{2} \left[\delta_{l_z, 0} h_{l_z}^-(p_n Z_{\max}) - S_{j'_z l'_z T_3^z, j_z l_z T_3^z} h_{l'_z}^+(p_n Z_{\max}) \right] C_{T_3 T_3^z, \frac{1}{2} - \frac{1}{2}}^{T0} \right. \\ &\quad \left. - \frac{i}{2} \sqrt{\frac{p'_p}{p_n}} S_{j'_z l'_z T_3^z, j_z l_z T_3^z} e^{2i\sigma_{l'_z}} u_{l'_z}^+(\eta, p'_p Z_{\max}) C_{T_3 T_3^z, \frac{1}{2} \frac{1}{2}}^{T0} \right). \end{aligned} \quad (13)$$

In the above equation, $p_n = \frac{3}{4}mv_{rel}$ is neutron's momentum of the n - ^3He channel, while $p_p = [p_n^2 + \frac{3m}{2\hbar^2} (B_{^3\text{H}} - B_{^3\text{He}})]^{1/2}$ is the proton's momentum of the p - ^3H recoils; $h_{l_z}^\pm$ are the spherical Hankel functions, and $u_{l'_z}^+$ is the outgoing Coulomb function of p - ^3H channel with the strength $\eta = \frac{4}{3}\alpha mc/(\hbar p_p)$. The functions $\{f_i^{\alpha a}(\mathbf{x}_i, \mathbf{y}_i)\}_{J_3 \equiv \frac{1}{2}, T_3 T_3^z}$ with $T_3^z = \pm \frac{1}{2}$ are the normalized Faddeev amplitudes for ^3He and ^3H , which we obtain by solving the corresponding $3N$ bound state problems.

The wave functions are dominated by the total isospin $\mathcal{T}=0$ state and, to describe the ^4He ground state, we can safely limit ourselves to the $\mathcal{T}=0$ state. However, in order to separate the n - ^3He and p - ^3H channels in the initial scattering state, we need to consider the admixture of total isospin $\mathcal{T}=1$ and $\mathcal{T}=2$ states; we therefore solve the scattering problem without constraining the total isospin value.

III. RESULTS

In this work we have performed rigorous FY calculations for five sets of nuclear Hamiltonians: Av18, I-N3LO, INOY, Av18+UIX and I-N3LO+UIX*. Here Av18 stands for the Argonne v18 potential [18], I-N3LO the chiral N³LO potential of Idaho group [19], INOY the semi-realistic configuration space potential that has been recently derived by Doleschall [20], and UIX is the tri-nucleon interaction derived at Urbana [21]. For the case of I-N3LO+UIX*, we have slightly change the parameters in UIX so as to reproduce exactly the triton binding energy, see Ref. [13] for details.

Our theory predictions for each of the model Hamiltonians used are listed in Table I. In addition to the *hen* cross section, the table also shows the binding energies of ^3H , ^3He

TABLE I: Theory predictions for the *hen* cross section, for five sets of the realistic nuclear Hamiltonians. We also list the values of binding energies (in units of MeV) for ${}^3\text{H}$, ${}^3\text{He}$ and ${}^4\text{He}$, the point-proton rms radius [23] r_{He4} (in fm) and the D -state probability $P_D({}^4\text{He})$ (in per cents) of the α -particle as well as the spin-triplet n - ${}^3\text{He}$ scattering length $a_{n\text{He3}}$ (in units of fm). The uncertainties attached to σ represent the variation of σ as the cutoff parameter Λ is varied in the range $\Lambda = (500 \sim 900)$ MeV.

	BE(${}^3\text{H}$)	BE(${}^3\text{He}$)	BE(${}^4\text{He}$)	r_{He4}	$P_D({}^4\text{He})$	$a_{n\text{He3}}$	σ [μb]
Av18	7.623	6.925	24.23	1.516	13.8	$3.43 - 0.0082i$	80.0 ± 12.2
I-N3LO	7.852	7.159	25.36	1.52	9.30	$3.56 - 0.0070i$	57.3 ± 7.9
INOY	8.483	7.720	29.08	1.377	5.95	$3.26 - 0.0058i$	34.4 ± 4.5
Av18+UIX	8.483	7.753	28.47	1.431	16.0	$3.23 - 0.0054i$	49.4 ± 8.5
I-N3LO+UIX*	8.482	7.737	28.12	1.475	10.9	$3.44 - 0.0055i$	44.4 ± 6.7
Exp.:	8.482	7.718	28.30	1.475(6)		$3.278(53) - 0.001(2)i$	$55 \pm 3, 54 \pm 6$

and ${}^4\text{He}$, the point-proton rms radius r_{He4} of ${}^4\text{He}$,^{#3} the D -state probability in ${}^4\text{He}$, and the spin-triplet n - ${}^3\text{He}$ scattering length $a_{n\text{He3}}$.

The bound state properties calculated in this work agree well with those obtained in the other calculations [24, 25, 26]. Theoretical calculations for n - ${}^3\text{He}$ scattering are much less established, but we have checked that our results agree within 2% with the momentum-space FY calculation of the Lisboa group [27], as well as with the RGM calculation carried out by Hofmann [28] for the Av18 and Av18+UIX Hamiltonians.

The table also indicates that the three-nucleon interactions (TNIs) play an important role in bringing the binding energies and the spin-triplet n - ${}^3\text{He}$ scattering length close to their respective experimental values. We remark that there is some uncertainty in the experimental value of the scattering length. The value listed in the table is due to an R -matrix analysis [29] based on the n - ${}^3\text{He}$ scattering data measured before the year 2002. Recently

^{#3} The point-proton rms radius r_{He4} is defined as $(r_{\text{He4}})^2 \equiv r_c^2({}^4\text{He}) - r_p^2 - r_n^2$, where $r_c({}^4\text{He})$ is the proton charge rms radius of ${}^4\text{He}$, r_p and r_n are the rms charge radius of the proton and neutron, respectively. See, for example, [22] for detailed explanation. With the ${}^4\text{He}$ proton charge radius $r_c({}^4\text{He}) = 1.681(4)$ fm obtained in a recent analysis [23] and with the 2008 PDG values for the proton and neutron rms radii, $r_p = 0.875(7)$ fm and $r_n^2 = -0.1161(22)$ fm², we arrived at $r_{\text{He4}} = 1.475(6)$ fm, which is about 1.4 % larger than the estimate given in [22].

new measurements of the coherent scattering length have been performed at NIST [30] and ILL [31], but the results of the two groups do not agree with each other and both of them are in disagreement with the old ILL measurement.

The D -state probability in the α -particle, $P_D(^4\text{He})$, is closely related to the tensor forces and shows strong model dependence. However this quantity is not an observable, and it turns out to be difficult to constrain this quantity by studying other processes that are sensitive to $P_D(^4\text{He})$ [32].

As can be seen from the Table, the calculated value of the *hen* cross section for Av18 (INOY) is too large (small), while the remaining three other Hamiltonians exhibit only a mild variation around the experimental values. The discrepancy of the Av18 prediction is not surprising, since this Hamiltonian lacks TNIs and fails to reproduce even the binding energies. While a full discussion on model dependence will be given in the next subsection, this is a good place to discuss a feature that is specific to INOY. Although the INOY potential is capable of reproducing the binding energies and rms radii of $A = 3$ system quite accurately [33], it is known to give too large binding energies and too small rms radii for $A \geq 4$ [34] and for nuclear matter [35]. Such a feature leads to the reduction of the overlap integration between n - ^3He and ^4He and hence to the underestimation of the *hen* cross section. The INOY potential also deviates from the Tjon-line (a line that correlates the $A = 3$ and $A = 4$ binding energies) rather severely, which also indicates that caution should be exercised in using INOY for the $A \geq 4$ systems.

For further discussion, we list in Table II the values of the *hen* matrix elements and the LECs evaluated at $\Lambda = 700$ MeV, for each of the nuclear Hamiltonians. The real part of $\mathcal{M}$ is written as the sum of 1B and 2B contributions, with the dependence on g_{4s} and g_{4v} also shown. Several remarks are in order here. Firstly, the 2B contributions turn out to be about two times as big as 1B but with an opposite sign, which is consistent with the observation made in [8, 11]. Secondly, there are substantial model-dependence even in the 1B sector, which might be traced to the fact that not all the adopted Hamiltonians do not accurately reproduce the ERPs that govern the long- r contributions of 1B. Finally, the inclusion of TNI(UIX) plays quite an important role in reducing the model dependence.

TABLE II: Theory predictions for $\mathcal{M}$ (in units of $\text{fm}^3/2$), obtained with $\Lambda = 700$ MeV. $\Re\mathcal{M}$ is written in the format of: (1B) + (2B w/o CT) + (g_{4s} -term) + (g_{4v} -term) = (total).

	g_{4s}	g_{4v}	$\Re\mathcal{M}$	$\Im\mathcal{M}$
Av18	0.3958	0.1947	$0.1531 - 0.3777 + 0.0237 g_{4s} - 0.0403 g_{4v} = -0.2231$	0.0249
I-N3LO	0.3919	2.7479	$0.1304 - 0.2248 + 0.0198 g_{4s} - 0.0371 g_{4v} = -0.1885$	0.0203
INOY	0.2313	0.8021	$0.1045 - 0.2035 + 0.0473 g_{4s} - 0.0725 g_{4v} = -0.1462$	0.0154
Av18+UIX	0.5810	-0.4615	$0.1518 - 0.3567 + 0.0205 g_{4s} - 0.0377 g_{4v} = -0.1756$	0.0179
I-N3LO+UIX*	0.5402	2.3249	$0.1305 - 0.2253 + 0.0175 g_{4s} - 0.0347 g_{4v} = -0.1661$	0.0183

A. Model-dependence

In understanding the observed model dependence, it seems informative to recall the results of our previous study [13] of the M1 properties of the $A \leq 3$ nuclei. It was found in [13] that the M1 matrix elements (MEs) of the $A=3$ systems are strongly correlated with the triton binding energy B_3 . Since B_3 governs the long-distance contributions to the MEs, the model-dependence (viz., variations in the MEs corresponding to the different nuclear Hamiltonians that give different values for B_3) cannot be cured by the renormalization procedure through the local (or short-ranged) operators. However, using the empirically established correlation curve between MEs and B_3 , we were able to calibrate the values of MEs by replacing the model-predicted B_3 to its experimental value, which resulted in an essentially model-independent theoretical prediction.

We expect that we can in principle adopt the same procedure for the *hen* process. Indeed we found that σ is almost proportional to the quantity

$$\zeta \equiv [q (a_{n\text{He}3}/r_{\text{He}4})^2]^{-2.75}. \quad (14)$$

We plot in Fig. 2 σ versus ζ for the five nuclear Hamiltonians under consideration. Fig. 2 suggests a strong correlation between σ and ζ . If we take this correlation seriously and consider the quantity $\tilde{\sigma}$ defined by

$$\tilde{\sigma} \equiv \frac{\zeta_{\text{exp}}}{\zeta} \sigma, \quad (15)$$

then $\tilde{\sigma}$ turns out to be almost model-independent:

$$\tilde{\sigma} = (54.8 \pm 8.4, 54.5 \pm 7.5, 53.9 \pm 7.1, 54.8 \pm 9.4, 56.4 \pm 8.5) \mu b \quad (16)$$




FIG. 2: The *hen* cross section σ (in the unit of μb) plotted against ζ/ζ_{exp} , where $\zeta = [q(a_{n\text{He}3}/r_{\text{He}4})^2]^{-2.75}$. For each Hamiltonian, the error bar denotes a range of variation for three different choices of the cutoff parameter: $\Lambda = 500$ MeV (bottom), 700 MeV (middle with a filled circle) and 900 MeV (top). The lines in red with the label “Exp” denote the experimental data, 54 ± 6 [5] and 55 ± 3 [6].

for Av18, I-N3LO, INOY, Av18+UIX and I-N3LO+UIX*, all of which are in perfect agreement with the data.

It is however not quite clear whether the correlation between σ and ζ is accidental or physical in nature. Furthermore, since *hen* is a four-body process that may involve a large number of ERPs, our numerical results that only cover five sets of Hamiltonians may not be sufficient to establish the correlation unambiguously. For example, a similar correlation behavior can be seen if we plot σ against $q^5 P_D^{2/3}$. (It is however not clear whether this is a separate correlation or just another way to express the correlation between σ and ζ , for P_D and $r_{\text{He}4}$ may not be independent of each other.) Without delving into the discussion of physics behind the correlation between σ and ζ , we take here the viewpoint that the σ calculated with a nuclear Hamiltonian that does not reproduce the relevant ERPs should be considered much less reliable than the σ obtained with a nuclear Hamiltonian that does reproduce the relevant ERPs. Based on this viewpoint, we adopt here the results

TABLE III: Cutoff-dependence of our prediction with the Av18+UIX wave function. The values of g_{4s} and g_{4v} are determined so as to reproduce the magnetic moments of the ^3H and ^3He nuclei. For $\Re\mathcal{M}$, its dependence on g_{4s} and g_{4v} are explicitly shown. $\mathcal{M}$ and σ are in units of $\text{fm}^{3/2}$ and μb , respectively.

Λ [MeV]	g_{4s}	g_{4v}	$\Re\mathcal{M}$	$\Im\mathcal{M}$	σ
500	0.8366	1.9068	$-0.0915 + 0.0246 g_{4s} - 0.0471 g_{4v} = -0.1608$	0.0166	40.9
600	0.6990	0.5886	$-0.1593 + 0.0231 g_{4s} - 0.0433 g_{4v} = -0.1686$	0.0173	44.9
700	0.5810	-0.4615	$-0.2049 + 0.0205 g_{4s} - 0.0377 g_{4v} = -0.1756$	0.0179	48.7
800	0.4517	-1.3622	$-0.2346 + 0.0176 g_{4s} - 0.0319 g_{4v} = -0.1832$	0.0186	52.9
900	0.3169	-2.2069	$-0.2547 + 0.0149 g_{4s} - 0.0265 g_{4v} = -0.1915$	0.0195	57.9

obtained with Av18+UIX and I-N3LO+UIX* as the most reliable predictions. For these two Hamiltonians, the calculated *hen* cross sections are:

$$\sigma = 49.4 \pm 8.5 \mu b \text{ (Av18 + UIX)}, \quad \sigma = 44.4 \pm 6.7 \mu b \text{ (I-N3LO+UIX*)}. \quad (17)$$

Both are lower by a few μb than the central value of the the experimental data but still in good agreement with the data within the error bars.

B. Cutoff dependence

We now turn our attention to the cutoff dependence. Table III shows to what extent our results for the Av18+UIX wave functions change when the cutoff parameter Λ is varied over a range $\Lambda = 500 \sim 900$ MeV. Table III indicates that the renormalization procedure of LECs (g_{4s} and g_{4v}) plays an essential role in removing the cutoff-dependence. As a way of quantifying its effectiveness, we define the quantity

$$R \equiv \frac{\mathcal{M}_{\text{total}}(\Lambda_2) - \mathcal{M}_{\text{total}}(\Lambda_1)}{\mathcal{M}_{\text{finite}}(\Lambda_2) - \mathcal{M}_{\text{finite}}(\Lambda_1)}, \quad (18)$$

where the subscript “finite” stands for “finite-range term contributions”, and it corresponds to a case where all the terms other than the contact term (CT) contributions are included. Perfect renormalization invariance would correspond to $R=0$. It turns out that $R_{\text{hen}} = 0.189$ for $(\Lambda_1, \Lambda_2) = (500, 900)$ MeV. Thus the renormalization procedure of LECs has

TABLE IV: Matrix elements for the Av18+UIX wave function with $\Lambda = 700$ MeV; the LEC corresponding to this case are: $g_{4s} = 0.581$ and $g_{4v} = -0.4615$.

	$\mu(^3\text{H})$	$\mu(^3\text{He})$	$\Re\mathcal{M}$	$\Im\mathcal{M}$
1B: LO	2.5727	-1.7632	0.0964	-0.0136
1B: RC	-0.0171	0.0037	0.0554	-0.0075
1B-total	2.5556	-1.7595	0.1518	-0.0211
2B: 1π (NLO)	0.2292	-0.2258	-0.1657	0.0195
2B: $1\pi C$ (N^3LO)	0.1578	-0.1289	-0.1465	0.0172
2B: 2π (N^3LO)	0.0419	-0.0408	-0.0445	0.0052
finite (total w/o CT)	2.9845	-2.1550	-0.2049	0.0208
	$0.0193 g_{4s}$	$0.0190 g_{4s}$	$0.0205 g_{4s}$	$-0.0014 g_{4s}$
	$+0.0363 g_{4v}$	$-0.0354 g_{4v}$	$-0.0377 g_{4v}$	$+0.0044 g_{4v}$
2B: CT (N^3LO)	$= -0.0055$	$= 0.0274$	$= 0.0293$	$= -0.0029$
Total	2.9790	-2.1276	-0.1756	0.0179

removed major part of cutoff-dependence, reducing the cutoff-dependence to about one-fifth of $\mathcal{M}_{\text{finite}}$. It is interesting to compare the above value of R_{hen} with the corresponding quantity R_{hep} obtained in a *hep* calculation in [3]. The *hep* calculation in [3] is based on the same MEEFT strategy (but used a different method for obtaining exact solutions to the nuclear Schroedinger equations). Ref. [3] reports $R_{\text{hep}} = 0.137$ for a slightly smaller window $(\Lambda_1, \Lambda_2) = (500, 800)$ MeV. Thus the previous *hep* calculation [3] is consistent with the *hen* calculation in the present work, and this consistency renders further support to the *hep* results in [3].

C. Details of $\mathcal{M}$ and chiral convergence

Table IV shows the individual contributions of the various 1B and 2B terms to $\mu(^3\text{H})$, $\mu(^3\text{He})$ and M_{hen} , calculated at $\Lambda = 700$ MeV for the Av18+UIX Hamiltonian. We can see that the 1B contribution to *hen* is highly suppressed due to the aforementioned orthogonality between the initial and final wave functions. The NLO contribution, which comes

from the *soft* one-pion-exchange, is also suppressed for the M1 channel, due to the accidental cancelation between the pion-pole and pion-seagull diagram contributions [13]. These suppression mechanisms make chiral convergence rather unclear. For example, one might worry about the fact that the $1\pi C$ and 2π contributions, both of which are N³LO, turn out to be comparable in size to the NLO 1π contribution. It should be noted, however, that most of the $1\pi C$ and 2π contributions are to be absorbed in the renormalization of the LECs, leaving very small effects on the net amplitudes. To demonstrate this, we define the *effective* matrix element, $\langle \mathcal{O} \rangle^{\text{effective}}$, of a given operator $\mathcal{O}$ by

$$\langle \mathcal{O} \rangle^{\text{effective}} \equiv \mathcal{M}_{\text{total}} - (\mathcal{M}_{\text{total}} \text{ but without } \langle \mathcal{O} \rangle). \quad (19)$$

Thus $\langle \mathcal{O} \rangle^{\text{effective}}$ represents a change in the net amplitude that would occur if we omit the operator $\mathcal{O}$. In evaluating the parenthesized quantity, the LECs should be readjusted so as to reproduce the experimental values of the $A = 3$ magnetic moments *without* $\langle \mathcal{O} \rangle$; because of this readjustment we should expect $\langle \mathcal{O} \rangle^{\text{effective}} \neq \langle \mathcal{O} \rangle$. We find

$$\begin{aligned} \langle 1\pi \rangle^{\text{effective}} &= 0.0749 - 0.0087i, & (g_{4s}, g_{4v}) &= (0.5223, 5.8848), \\ \langle 1\pi C \rangle^{\text{effective}} &= -0.0093 + 0.0004i, & (g_{4s}, g_{4v}) &= (1.2433, 3.53455), \\ \langle 2\pi \rangle^{\text{effective}} &= -0.0010 + 0.0001i, & (g_{4s}, g_{4v}) &= (0.5843, 0.6921), \end{aligned} \quad (20)$$

where we have also listed the corresponding values of the LECs, which should be compared with $(g_{4s}, g_{4v}) = (0.581, -0.4615)$ that corresponds to the full calculation up to N³LO. Eq.(20) demonstrates that the *effective* contributions of $1\pi C$ and 2π are very small, only about 6 % and 2 %, respectively, relative to the values one would naively expect. This is in sharp contrast with the NLO soft one-pion-exchange, whose contribution cannot be absorbed in the LECs. A rigorous proof of chiral convergence would require a calculation that goes one order higher than considered in the present work; *i.e.*, we need to go up to N⁴LO.

IV. DISCUSSION AND CONCLUSION

There can also be a “completely” consistent approach where both nuclear Hamiltonians and transition operators are obtained in the same EFT framework. This approach requires much more involved calculations than MEEFT, but recent significant progress in constructing EFT Hamiltonians makes it more attractive. Also available is a pionless EFT

approach [37] where the matrix elements are evaluated perturbatively, but, unless it is capable of reproducing all the relevant ERPs of the nuclear systems under consideration, its usefulness is limited.

In this work we have performed an *ab initio* parameter-free calculation for the *hen* cross section, with the EM currents that have been derived from HBChPT up to N³LO. Exact nuclear wave functions have been obtained by solving the Faddeev-Yakubovskii equations for realistic nuclear Hamiltonians. The calculated value of σ shows a high degree of stability as the cutoff parameter Λ is varied over a wide range, $\Lambda = (500 \sim 900)$ MeV, and we obtain as the best estimate $\sigma = 49.4 \pm 8.5 \mu b$ for Av18+UIX and $44.4 \pm 6.7 \mu b$ for I-N³LO+UIX*. These values are in good agreement with the data, 54 ± 6 [5] and 55 ± 3 [6].

The successful application of MEEFT to *hen* renders strong support to the previous MEEFT calculation for *hep* in Ref. [3]; furthermore, it demonstrates the basic soundness of the MEEFT strategy in general. The present treatment is open to several improvements such as: the inclusion of the next order in chiral perturbation, in particular the incorporation of the three-nucleon currents; a more stringent control of mismatch in the chiral counting between SNPA and a formally accurate chiral expansion that enters in the currents; a better understanding of the role that the counter terms play in the renormalization group property, etc. It is reasonable, however, to expect that the effects of these improvements are essentially accommodated in the above-quoted error estimate based on the cutoff dependence. A robust estimation of the *hep* S-factor has been a long-standing challenging problem in nuclear physics [36]. We believe that our MEEFT calculations of *hep* and *hen* have solved this problem to a satisfactory degree. Short of doing a fully formally consistent EFT calculation, which is at present out of our reach, MEEFT seems to be the best currently available approach.

Acknowledgement

The work of TSP is supported by the Korea Science and Engineering Foundation (KOSEF) Basic Research Program with the grant No. R01-2006-10912-0, by the KOSEF grant funded by the Korea Government (MEST) (No. M20608520001-08B0852-00110) and by the US National Science Foundation, Grant No. PHY-0758114.

The numerical calculations have been performed at IDRIS (CNRS, France). We thank

the staff members of the IDRIS computer center for their constant help. We are deeply obliged to Professors Mannque Rho and Kuniharu Kubodera for kindly agreeing to read the manuscript prior to its publication and giving us important comments.

-
- [1] K. Kubodera and T.-S. Park, Ann. Rev. Nucl. Part. Sci. **54** (2004) 19.
 - [2] INT workshop on “Solar Fusion Cross Sections for the pp chain and CNO cycle”, Institute for Nuclear Theory, Seattle, USA, January 21-23, 2009.
 - [3] T.-S. Park, L.E. Marcucci, R. Schiavilla, M. Viviani, A. Kievsky, S. Rosati, K. Kubodera, D.-P. Min and M. Rho, Phys. Rev. C **67** (2003) 055206.
 - [4] G.E. Brown and M. Rho, Phys. Rept. **396** (2004) 1.
 - [5] F.L.H. Wolfs, S.J. Freedman, J.E. Nelson, M.S. Dewey and G.L. Greene, Phys. Rev. Lett. **63** (1989) 2721.
 - [6] R. Wervelman, K. Abrahams, H. Postma, J.G.L. Booten and A.G.M. Van Hees, Nucl. Phys. **A526** (1991) 265.
 - [7] I.S. Towner and F.C. Khanna, Nucl. Phys. **A356** (1981) 445.
 - [8] J. Carlson, D.O. Riska, R. Schiavilla and R.B. Wiringa, Phys. Rev. C **42** (1990) 830.
 - [9] R. Schiavilla, R.B. Wiringa, V.R. Pandharipande and J. Carlson, Phys. Rev. C **45** (1992) 2628.
 - [10] M. Chemtob and M. Rho, Nucl. Phys. **A163** (1971) 1.
 - [11] Y.-H. Song and T.-S. Park, arXiv:nucl-th/0311055.
 - [12] Y.-H. Song, R. Lazauskas and T.-S. Park, Phys. Lett. B **656** (2007) 174.
 - [13] Y.-H. Song, R. Lazauskas and T.-S. Park, nucl-th/0812.3834.
 - [14] L.D. Faddeev, Zh. Eksp. Teor. Fiz. **39** (1960) 1459 (Wiley & Sons Inc., 1972) [Sov. Phys. JETP **12** (1961) 1014].
 - [15] O.A. Yakubowsky, Sov. J. Nucl. Phys. **5** (1967) 937.
 - [16] R. Lazauskas, Ph.D. Thesis, Université Joseph Fourier, Grenoble (2003); <http://tel.ccsd.cnrs.fr/documents/archives0/00/00/41/78/>.
 - [17] R. Lazauskas, arXiv:0808.1650 [nucl-th]; Few-Body Syst, DOI 10.1007/s00601-008-0006-3.
 - [18] R.B. Wiringa, V.G.J. Stoks, R. Schiavilla, Phys. Rev. C **51** (1995) 38.
 - [19] D.R. Entem and R. Machleidt, Phys. Rev. C **68** (2003) 041001(R).

- [20] P. Doleschall, I. Borbély, Z. Papp, W. Plessas, Phys. Rev. C **67** (2003) 064005.
- [21] B.S. Pudliner, V.R. Pandharipande, J. Carlson and R.B. Wiringa, Phys. Rev. Lett. **74** (1995) 4396.
- [22] E. Caurier and P. Navrátil, Phys. Rev. C **73** (2006) 021302(R).
- [23] I. Sick, Phys. Rev. C **77** (2008) 041302(R).
- [24] A. Deltuva and A.C. Fonseca, Phys. Rev. C **75** (2007) 014005.
- [25] A. Kievsky, S. Rosati, M. Viviani, L.E. Marcucci and L. Girlanda, J. Phys. G **35** (2008) 063101.
- [26] A. Nogga, H. Kamada and W. Glöckle, Phys. Rev. Lett. **85** (2000) 944.
- [27] A. Deltuva, private communication.
- [28] H.M. Hofmann and G.M. Hale, arXiv:nucl-th/0211008.
- [29] M.G. Huber, M. Arif, T.C. Black, W.C. Chen, T.R. Gentile, D.S. Hussey, D. Pushin, F.E. Wietfeldt and L. Yang, arXiv:0809.2248 [nucl-ex].
- [30] P.R. Huffman, D.L. Jacobson, K. Schoen, M. Arif, T.C. Black, W.M. Snow and S.A. Werner, Phys. Rev. C **70** (2004) 014004.
- [31] W. Ketter, W. Heil, G. Badurek, M. Baron, E. Jericha, R. Loidl and H. Rauch, Eur. Phys. J. A **27** (2006) 243.
- [32] A. Arriaga, V.R. Pandharipande and R. Schiavilla, Phys. Rev. C **43** (1991) 983.
- [33] R. Lazauskas and J. Carbonell, Phys. Rev. C **70** (2004) 044002.
- [34] C. Forssen, E. Caurier and P. Navratil, Phys. Rev. C **79** (2009) 021303(R).
- [35] M. Baldo and C. Maieron, Phys. Rev. C **72** (2005) 034005.
- [36] J.N. Bahcall, M.H. Pinsonneault and S. Basu, Astrophys. J. **555** (2001) 990.
- [37] P.F. Bedaque and U. van Kolck, Ann. Rev. Nucl. Part. Sci. **52** (2002) 339 and references therein.